

12 .
 1/3
 96 %
 24 .
 100
 96 %.
 40 ° ,
 [8].
 « »
 () .
 0,15 % 1:10
 1 .
 96 %
 96 %
 () .
 7 % 1:10
 12
 24 .
 25
 96 %
 96 % [1, 2, 5].
 L.
 2009-2012 . .
 ()
 70 %
 1:10 3
 1:10

	, %*			
	9,80±2,91	2,39±0,84	8,282±0,420	8,270±1,380
	8,080±1,055	1,95±0,73	13,254±0,530	4,580±0,628
	< 1,0	< 1,0	22,012±1,640	8,050±0,981
	7,38±0,94	1,76±0,68	14,84±1,32	5,19±0,38

* : = 5.

(1 /). -
 (6:4:3)
 105 ° 15 .
 [6, 7, 9].
 ()
 7 10 %.

1.
 2.
 3.

1. // . 79-83.
 - 2008. - 2. - . 135-136.
 2. // . 2001. - 556 .
 - 84 .
 3. // . - 1970.
 : , . , 1990. - 333 .
 4. // . - 2005. - N 4. - . 16-17.
 5. // . - 2005. - 6. -

6. / « -
 , 2001. - 556 .
 7. /
 . - 2007. - 2. - . 38-39.
 8. Brent L. Ridley *Pectins: structure, biosynthesis, and oligogalacturonide-related signaling*. /L. Brent Ridley, A. Malcolm [et al.] // *Phytochem.* — 2001. - 57. - . 929-967.
 9. *Thin-layer chromatography: reagents and detection* / H. Jork, W. Funk, W. Fischer, H. Wimmer. - Weinheim; New York: VCH. - 1990. - Vol. 15. - 497p.
 05.03.2013

: 633.854.78 +635.24] 577.114_

N. P. Maksyutina, O. I. Yemelianova, Y. A. Tsymbalista
STUDY OF COMPOSITION OF POLYSACCHARIDES
HELIANTHUS TUBEROSUS AND HELIANTHUS ANNUUS

Keywords: polysaccharides, *Helianthus tuberosus*, *Helianthus annuus*

The article presents data on the allocation of research and polysaccharide fractions from tubers and leaves of *Helianthus tuberosus* and roots, leaves *Helianthus annuus*. Soluble fraction was obtained polysaccharides, pectin, hemicellulose A and B and installed quantitative composition.