

## CONTENTS

---

Behaviour of topological defects of optical indicatrix orientation in cubic, hexagonal, trigonal and tetragonal crystals under conically distributed electric field. 2. The electric field and the optical beam parallel to the principal crystallographic directions <i>Vasylykiv Yu., Kryvyy T., Skab I. and Vlokh R.</i>	1
Wide-slit diffraction and wavefront diagnostics of optical-vortex beams <i>Bekshaev A. Ya., Kurka I. A., Mohammed K. A. and Slobodeniuk I. I.</i>	17
The influence of structural defects on the optical properties of synthetic opals <i>Yevchik A., Moiseyenko V. and Dergachov M.</i>	24
A Q-switched fibre laser operating in the 2 $\mu\text{m}$ region based on nonlinear polarization rotation technique <i>Azooz S. M., Harun S. W., Ahmad H., Halder A., Paul M. C., Das S. and Bhadra S. K.</i>	32
Anisotropy of acoustooptic figure of merit for $\text{TeO}_2$ crystals. 2. Anisotropic diffraction <i>Mys O., Kostyrko M., Krupych O. and Vlokh R.</i>	38
Measurements of ozone absorption cross section with ratiometric and non-ratiometric methods <i>Tay Ching En Marcus, Mohd Haniff Ibrahim and Nor Hafizah Ngajikin</i>	61

---