

UDC 57.085.25:541.1/.49



Rybachuk O. A.<sup>1,2,4</sup>, Levin R. E.<sup>1,3</sup>, Kyryk V. M.<sup>2</sup>, Susarova D. K.<sup>3</sup>, Tsupykov O. M.<sup>1,2,4</sup>, Smozhanik E. G.<sup>1,4</sup>, Butenko G. M.<sup>2</sup>, Skibo G. G.<sup>1,2,4</sup>, Troshin P. A.<sup>3</sup>, Pivneva T. A.<sup>1,2,4</sup>

<sup>1</sup>Bogomoletz Institute of Physiology NAS Ukraine, Kyiv, Ukraine

<sup>2</sup>State Institute of Genetic and Regenerative Medicine NAMS Ukraine, Kyiv, Ukraine

<sup>3</sup>Institute of Problems of Chemical Physics, Russian Academy of Sciences, Chernogolovka, Russia

<sup>4</sup>State Key Laboratory, Kyiv, Ukraine

e-mail: oks-ribachuk@yandex.ru

# EFFECT OF A WATER SOLUBLE DERIVATIVE OF FULLERENE C<sub>60</sub> ON THE FEATURES NEURAL PROGENITOR CELLS *IN VITRO*

## ABSTRACT

We studied the effect of a water soluble derivative of fullerene C<sub>60</sub> on the behavior of cultured neural stem/progenitor cells. Addition of 20 nM of metal fullerlenolate C<sub>60</sub> (NaFL) into the cell culture increased the population of the cells almost twice in comparison with the control and also suppressed the formation of neurospheres. The obtained data allow us to suggest that NaFL has a positive effect on the proliferative activity of neural progenitors. The water-soluble fullerene nanostructures such as NaFL promoting the proliferation of neural stem cells might have numerous beneficent applications in cell biology and biotechnology.

**KEYWORDS:** neural stem cells, NaFL, cell proliferation, neurospheres, FGF-2, immunocytochemical staining, electron microscopy.

One of the perspective areas in cell and combined gene-cell therapy is the selection, identification of stem cells, and obtaining of stem clones, especially pluripotent cells; and their subsequent differentiation into specialized cells [1].

Neural stem cells (NSC) can differentiate into neurons, astrocytes and oligodendrocytes, providing required amount of cells in the brain during its development and pathology. The study of NSC properties in culture conditions plays a huge role in the research of central nervous system (CNS) diseases, neurons development, its differentiation and regeneration.

Detection of neural stem cells and their identification in the brain of embryos and adults transformed the conception about the CNS plasticity. To replace the postulate that the neural cells do not renew and divide, there appeared the statement that the functions of damaged neurons in the adults are renewed by NSC. At present there has already been collected quite a large number of experimental data on the possible use of the NSC in various neurodegenerative diseases.

At present, the researchers' attention has been attracted by the problem of isolation, multiplication, long-term storage of the NSC, and the study of their differentiation during the cultivation and after transplantation into the brain. In the adult stem cell number is much less than in the embryonic and early postnatal period. Therefore, there is not enough donor material for transplantation. Their ability to proliferate *in vitro* is used to increase the amount of stem cells.

The main advantage of cultivated cells is a capability for *in vivo* monitoring of cells using a microscope. Essential is that, working with cell cultures in the experiment, healthy cells only are used, and they remain

viable throughout the experiment. It is possible to ensure that, testing cell culture periodically. Moreover, it is easy to estimate the relative number of viable cells.

In the culture, the NSC maintained as self-renewing in clones, immature multipotent cells with a set of specific phenotypic markers. After proliferation blocking, they differentiate into neurons, oligodendrocytes and astroglia [2]. *In vitro* cultivated NSC is a widest source of cells for transplantation, drug screening, gene therapy and neuronal development studies [3-6].

At present there is clearly identified regulatory and selective effect of some growth factors on NSC properties both, *in vivo* and *in vitro* [7]. However, despite the early successes in the study of the NSC, there are still many issues and problems related to the study of these cell properties, development of methods for their isolation, growth, and differentiation.

At present, nanotechnology is considered the most promising of all the technologies, because there is a qualitative leap from the manipulation of the material to the manipulation of individual atoms [8, 9]. One of perspective directions is the use of carbon nanostructures — derivatives of fullerene (C<sub>60</sub>). Fullerene — is a molecular compound that belongs to the class of the allotropic forms of carbon, which are convex closed polyhedral composed of an even number of three-coordinated carbon atoms. In molecules of fullerene, carbon atoms are positioned at corners of regular hexagons and pentagons. The surface of a sphere or ellipsoid is composed of them.

Recent studies point to an antioxidant [10-18], neuroprotective [19-23] and nootropic [24] properties of C60 fullerene derivatives [10-24]. The authors also report about radioprotective [17], oncoprotective [25, 26],

antiallergic [27], antibacterial, and antiviral [28-36] properties of water-soluble  $C_{60}$  fullerene derivatives.

Despite such an impressive list of biological effects of fullerenes, we have to recognize that the mechanisms of many of them still have not been decrypted. There is a lack of knowledge of investigated nanoparticles pharmacodynamics and pharmacokinetics. There are not schemes of treatment for pulmonary, oncological, neurodegenerative and other diseases by fullerenes. Therefore, this trend requires more detailed research.

In this context, the aim of this research is to study the properties of the NSC by the action of water-soluble metal fullereneolate  $C_{60}$  (NaFL).

## MATERIALS AND METHODS

### ISOLATION OF NEURAL PROGENITORS AND ANALYZE OF THEIR VIABILITY

Under sterile conditions hippocampuses from the brain of FVB mice (17-18 days of embryonic development) were isolated. Then they were mechanically dissociated in a Neurobasal medium (Gibco, USA). Received cell suspension was passed through a 40  $\mu$ m nylon cell strainer (Falcon, USA). Purified fraction of NSC was prepared by centrifugation of cell suspension in a density gradient (22% Percoll). The percentage of viable cells in the suspension was determined by flow cytometry, using a cell sorter FACSAria (Becton Dickinson, USA), after incubation of cells with 7-amino-actinomycin D (7-AAD).

### CULTIVATION OF NEURAL PROGENITORS SUPPLEMENTED WITH NAFL

Neural progenitors were plated on Matrigel-coated glasses in 24-well plate ( $8 \cdot 10^4$  cells in each well) and cultivated in a  $CO_2$  incubator ( $t = +37^\circ C$ , 5%  $CO_2$ ). The first 2 days the cells were cultured in a standard Neurobasal medium (Gibco, USA) supplemented with FGF-2, as a growth factor that affects the proliferation of neural progenitors. In the absence of FGF-2 in the culture medium, the cells begin to differentiate spontaneously or form neurospheres.

Neural progenitors were cultivated under different conditions:

- 1 – standard conditions, FGF-2 is present in the culture medium;
- 2 – standard conditions + 20 nM NaFL;
- 3 – FGF-2 is not added in the culture medium;
- 4 – culture medium without FGF-2 + 20 nM NaFL.

The cells have been cultivated for 9 days after plating, 7 days in the presence of NaFL.

### ANALYZE OF PROLIFERATIVE ACTIVITY OF NEURAL PROGENITORS, STATISTICAL PROCESSING OF THE DATA

It is known that the proliferative (mitotic) activity of NSC depends on the presence of certain growth factors (FGF-2, EGF, NGF, BDNF, NT-3, NT-4) in the medium [7, 37-39]. Microenvironmental factors, in most cases, determine the fate of neural progenitors.

Proliferative activity of neural progenitors isolated from the hippocampus of 17-18-day mouse embryos was observed, using an inverted light microscope (Zeiss Telavar 31, Germany). On the 9th day after plating, the maximum cell density was observed in the wells to which 20 nM concentration of metallofullerene  $C_{60}$  was added.

Then, the cell culture was fixed with 4% paraformaldehyde for 30 minutes, washed in 0,1 M phosphate buffered saline (PBS) and enclosed into balsam.

Quantitative analysis of neural stem cells and neurospheres was performed, using an inverted microscope with a relief phase contrast (Olympus IX 71, Japan).

### IMMUNOCYTOCHEMICAL STAINING OF CULTIVATED NSC

To identify the neural progenitors and potential way of their differentiation there was used a double immunocytochemical staining for Nestin,  $\beta$ -tubulin III. Nestin is a specific marker of neuronal stem progenitors, post-mitotic neurons, and early neuroblasts [40].  $\beta$ -tubulin III is a specific marker of mature neuronal cytoskeleton.

Cell cultures were fixed with 4% paraformaldehyde for 30 minutes, washed with a 0,1 M PBS, and treated with 0,3% TritonX-100, 0,5% bovine serum albumin. During the following 24 hours they were incubated in a mixture of murine monoclonal Nestin (1:300), rabbit polyclonal  $\beta$ -tubulin III (1:750) of primary antibodies.

After washing off 0,1 M PBS, cell cultures were treated with a mixture of secondary anti-mouse Alexa Fluor 488 (Invitrogen, USA) 1:1000 and anti-rabbit Alexa Fluor 647 (Invitrogen, USA) 1:1000 antibodies for 1 hour.

Then the cells were washed in PBS and enclosed in the medium FluMount.

Fluorescence was visualized using selective filters (488 nm for Alexa Fluor 488, 647 nm for Alexa Fluor 647). This method allowed us to detect the presence of proliferating neural progenitors, and observe the morphological characteristics of neurospheres. Optical studies were performed using a laser scanning confocal microscope FluView™ FV1000 (Olympus, Japan).

### ELECTRON MICROSCOPIC RESEARCH OF CULTIVATED HIPPOCAMPAL SC

Cultures of neural progenitors were fixed with 2,5% glutaraldehyde solution and 2% paraformaldehyde for 1 hour at room temperature. Subsequently, cell cultures were washed with 0,1 M PBS, and fixed in 1%  $OsO_4$  solution for 1 hour; dehydrated in ethanol (30%, 50%, 70%, 80%, 96% absolute alcohol) and in 100 % acetone. Culture cells were enclosed in epoxy resins. The final step was a polymerization at 56 °C.

SC preparations, enclosed in a resin, were attached to the flat surface of the blocks with an adhesive «Cyanopan» (Poland). Ultrathin sections (500 Å) were cut on a microtome (LKB Bromma 8800, Sweden). They were contrasted with solutions of uranylacetate and lead citrate. Morphological characteristics of neural progenitors were evaluated with a transmission electron microscope JEM-100 CX (JEOL, Japan).

### STATISTICAL PROCESSING OF THE DATA

Statistical processing was carried out with the Student t-test, significance of the difference was taken at  $p < 0.001$ .

## RESULTS AND DISCUSSION

Using flow cytometry method there was determined that the percentage of viable cells after incubation with 7-amino-actinomycin (7-AAD) ranged from 89,8% to 93,5%.

Neural progenitors have been cultivated for 9 days after planting, and with NaFL — for 7 days with and without FGF-2.

It is established that FGF-2 causes proliferation of multipotent cells, which are the progenitors of neurons and glia in the subventricular zone of the lateral ventricles of the brain. FGF-2 together with NGF (neuronal growth factor) can maintain the generation and viability of neurons, which are cell derivatives of the subventricular zone. This way they promote neurogenesis in the adult human brain, as due to its natural reserves, or by implantation of progenitor cells [41]. In pharmacokinetics study of FGF-2 has been established that, after subcutaneous injection of 5 ng/g of body weight dose into rats, its concentration in blood and brain liquor rapidly increases, indicating

its free passing through haemathoencephalic barrier. In adult animals it increased the amount of *NSC* mitoses in the subventricular zone and olfactory tracts, regulating current neurogenesis with particular humoral method [38]. Therefore, *FGF-2* performs the complex regulation of the de novo generation of brain cells [7]. During cultivation of multipotent stem cells, growth factor causes intense proliferation; and its removal from the medium induces their differentiation into neurons and glia. [39]

In our case, the neural progenitors, which were cultured with the addition of *FGF-2*, remained proliferative potential without formation of neurospheres (Fig. 1, A). In the wells without *FGF-2* a small amount of neural progenitors was remained. Also there was observed an appearance of neurospheres. That indicates the loss of neural progenitor's proliferative potential (Fig. 1, B).

Quantitative microscopic analysis of *NSC* under various conditions of cultivation showed that in the control, where in the culture medium *FGF-2* was present,  $543 \pm 61$  cells per  $\text{mm}^2$  appeared (Fig. 1, A; Fig. 2); at the same time, the amount of *NSC*, adding 20 nM concentration of *NaFL*, remained practically unchanged —  $530 \pm 66$  cells per  $\text{mm}^2$  (Fig. 1, B; Fig. 2).

In those wells, where the cells were cultured without *FGF-2*, the appearance of neurospheres was observed, the number of neural progenitors was  $200 \pm 77$  cells per  $\text{mm}^2$  ( $p < 0.001$ ) (Fig. 1, C; Fig. 2). By adding 20 nM concentration of *NaFL* (missing the growth factor in the culture medium), we observed a significant increase of *NSC* —  $426 \pm 101$  cells per  $\text{mm}^2$  ( $p < 0.001$ ) (Fig. 1, D; Fig. 2).

These data indicate that in the presence of 20 nM concentration of *NaFL* in the culture medium, but without growth factor, the number of neural progenitors is about 2 times higher than in control wells.

It can be assumed that *NaFL* increases the proliferative potential of neural progenitors by still unknown mechanism of action.

Method of immunocytochemical staining of cell culture has given us the ability to visualize neural progenitors and neurospheres in different types of cultivation. Neural progenitors at cultivation in the usual medium and the medium with 20 nM concentration of *NaFL* were

morphologically identical and showed practically the same proliferation activity (Fig. 3, A-E).

This result allows us to assume that 20 nM concentration of *NaFL* in this version of cultivation is not toxic, but did not affect the activity of *NSC* proliferation.


In immunocytochemical analysis of *NSC* cell culture, which was cultured in medium without *FGF-2*, we observed the formation of neurospheres, as well as a small number of neural progenitors. Currently it is not considered something exceptional to obtain the culture growth by type formation of neurospheres. This opens new possibilities for a thorough examination of cells within a neurosphere, and their subtyping [42]. It is obvious that, during the formation of neurosphere, more complex functions of undifferentiated neural cells appear than at cultivation in adherent mediums. Because the genesis of such a unique multi-cellular structures as a neurosphere requires the *NSC* ability to complex cellular interactions [43].

It is remarkable that today the *NSC* is the only type of stem cells, which possibility of such self-sustaining colony formation in promitotic environment is described [7].


Our study demonstrated that the addition of 20 nM concentration of *NaFL* in culture medium, which contained no *FGF-2*, increases the number of neural progenitors, and decreases the number of neurospheres (Fig. 4, D-E). This result confirms that the 20 nM concentration of *NaFL* increases the proliferative potential of *NSC*.

The electron-microscopic analysis showed a mitotic activity during the cultivation of neural progenitors in the medium, which contained *FGF-2*. This is characteristic to this cell type (Fig. 5, A). But, when we added *NaFL* into the culture medium, by the 7th day it was found not only outside the cells, but was localized near the cell membrane, in the membrane itself, and in the cytoplasm (Fig. 5, B-E).

Electron-microscopic investigations have shown that metal fullereneolol  $C_{60}$  may not only be present in the solution, but also penetrates into the cells directly across the cell membrane without damaging it; and localizes itself in the cytoplasm of cells, though this substance did not show any specific tropism for neural progenitors organelles.




**Fig. 1.** Neural stem cells and neurospheres, 7 days in culture. A – control (*NSC* + *FGF-2*); B – *NSC* + *FGF-2* with the addition of 20 nM concentration *NaFL*; C – *NSC* without *FGF-2* - neurospheres control; D – *NSC* cultivated without *FGF-2* + 20 nM *NaFL*.




**Fig. 2.** Quantitative analysis of *NSC* under different culture conditions, 7 days: 1 – standard conditions, *FGF-2* is present in the cultivation medium; 2 – standard conditions + 20nM *NaFL*; 3 – the cultivation medium without *FGF-2*; 4 – cultivation medium without *FGF-2* + 20 nM *NaFL*.






**Fig. 3.** The immunocytochemical staining of neural progenitors cultured in medium containing *FGF-2*, 7 days (x40):

**A** – control, *Nestin*-positive cells (green);  
**B** – control of  $\beta$ -*tubulin III*-positive cells (red);  
**C** – *Nestin* +  $\beta$ -*tubulin III*, control,  
**D** – NSC + NaFL, *Nestin*-positive cells (green),  
**E** – NSC + NaFL  $\beta$ -*tubulin III*-positive cells (red), and  
**F** – *Nestin* +  $\beta$ -*tubulin III*, NSC + NaFL.



**Fig. 4.** The immunocytochemical staining of neural progenitors and neurospheres cultured in medium without *FGF-2* 7 days (x20).

**A** – control (*Nestin*, green);  
**B** – control ( $\beta$ -*tubulin III*, red);  
**C** – *Nestin* +  $\beta$ -*tubulin III*, control;  
**D** – NSC + NaFL (*Nestin*, green);  
**E** – NSC + NaFL ( $\beta$ -*tubulin III*, red);  
**F** – *Nestin* +  $\beta$ -*tubulin III*, NSC + NaFL.



**Fig. 5.** Electron-microscopic images of cultured NSC:

**A** – dividing cell, control;

**B** – localization of  $MC60(OH)_x$  in the membrane and in the cytoplasm of cells;

**C** – localization of  $NaFL$  outside the cell processes;

**D** – localization of  $NaFL$  in cytoplasm;

**E** – localization  $NaFL$  near the membrane and in the cytoplasm.

Scale bar A — 0,7  $\mu\text{m}$ , B, C, D — 0,3  $\mu\text{m}$ .

## CONCLUSIONS

IT HAS BEEN SHOWN THAT UNDER THE CONDITIONS IN VITRO THE NUMBER OF CULTURED NSC, ADDING 20 NM CONCENTRATION OF METAL FULLERENOLAT  $C_{60}$ , INCREASES ALMOST 2 TIMES, COMPARED WITH THE CONTROL. THEIR PROLIFERATIVE ACTIVITY REMAINS THROUGHOUT THE PERIOD OF CULTIVATION, AND ALSO THE FORMATION OF NEUROSPHERES DECREASES. ULTRASTRUCTURAL ANALYSIS INDICATED THAT  $NAFL$  ENTERS THE CELL DIRECTLY THROUGH THE CELL MEMBRANE, WITHOUT DAMAGING IT; AND LOCALIZES IN THE CELL CYTOPLASM.

THUS, THE RESULTS OF THIS STUDY LEAD TO THE HYPOTHESIS THAT THE  $NAFL$  AFFECTING THE PROLIFERATION OF NEURAL STEM CELLS, MAY BE PERSPECTIVE FOR ITS USE IN CELL BIOLOGY AND BIOTECHNOLOGY.

## REFERENCES

1. Репин В. С. Эмбриональная стволовая клетка: от фундаментальных исследований в клинику // Пат. физиол. и эксперим. терапия. – 2004 – №2. – С. 3-8.
2. Price J., Williams B. Neural stem cells // Curr. Opin. Neurobiol. – 2001. – **11**, № 5. – P. 564-87.
3. Goldman S. Stem and progenitor cell-based therapy of the human central nervous system // Nat. Biotechnol. – 2005. – **23**, № 7. – P. 862–871.
4. Ourednik V., Ourednik J., Xu Y. et al. Cross-talk between stem cells and the dysfunctional brain is facilitated by manipulating the niche: evidence from an adhesion molecule // Stem Cells. – 2009. – **27**, №11. – P. 2846–2856.
5. Prestoz L., Relvas J., Hopkins K. et al. Association between integrin-dependent migration capacity of neural stem cells in vitro and anatomical repair following transplantation // Mol. Cell Neurosci. – 2001. – **18**, №5. – P. 473–484.
6. Svendsen C., Smith A. New prospects for human stem cell therapy in the nervous system // Trends Neurosci. – 1999. – **22**, №8. – P. 357–364.
7. Gage F. Mammalian neural stem cell // Science. – 2000. – **287**, № 5457. – P. 1433-1438.
8. Бурлакова Е. Б., Конрадов А. А., Мальцева Е. Л. Действие сверхмалых доз биологически активных веществ и низкоинтенсивных физических факторов // Химическая физика. – 2003 – **22**, № 2. – С 21–40.
9. Хадарцев А. А., Туктамышев И. Ш. Шунгит в медицине. Информационный материал. – Тула, 2005. – С. 5–6.
10. Tikhomirov A. A., Andrievsky G. V. Chronic alcoholization-induced damage to astroglia and intensification of lipid peroxidation in the rat brain protector effect of hydrated form of fullerene  $C_{60}$  // Neurophysiology. – 2007. – **39**, № 2. – P. 119–125.
11. Скулачев В. П. H2O2-сенсоры легких и кровеносных сосудов и их роль в антиоксидантной защите организма // Пульмонология. – 2000. – № 2. – С 6–9.
12. Wang L., Tai L., Lee D. et al.  $C_{60}$  and Water-Soluble fullerene derivatives as antioxidants against radical-initiated lipid peroxidation // J. Med. Chem. – 1999. – **4**, № 42(22). – P. 4614–4620.
13. Bensasson R., Bretteich M., Frederiksen J. et al. Reactions of (aq),  $CO_2^-$ ,  $HO^{\cdot}$ ,  $O_2^-$  and  $O_2$  with a dendro[60]fullerene and  $C_{60}[C(COOH)_2]_n$  ( $n = 52-6$ ) // Free Radical Biology and Medicine. – 2000. – **29**, №1. – P. 26–33.
14. Ali S., Hardt J., Quick K. et al. Biologically effective fullerene ( $C_{60}$ ) derivative with superoxide dismutase mimetic properties // Free Radical Biology and Medicine. – 2004. – **37**, № 8. – P. 1191–1202.
15. Chueh S., Lai M., Lee M. et al. Decrease of free radical level in organ perfusate by a novel water-soluble carbon-sixty, hexa (sulfobutyl) fullerenes // Transplant Proc. – 1999. – **31**, №5. – P. 1976–1977.

16. Avdeev M., Khokhryakov A., Tropin T. Structural features of molecular-colloidal solutions of C<sub>60</sub> fullerenes in water by small-angle neutron scattering // *Langmuir*. – 2004. – **20**, № 11. – P. 4363–4368.
17. Mchedlov-Petrossyan N., Klochkov V., Andrievsky G. Colloidal dispersions of fullerene C<sub>60</sub> in water: some properties and regularities of coagulation by electrolytes // *J. Chem. Soc., Faraday Trans.* – 1997. – **93**. – P. 4343–4346.
18. Huang S., Mashino T., Mochizuki M. et al. Effect of hexasulfobutylated C<sub>60</sub> on the isolated aortic ring of guinea pig // *Pharmacology*. – 2000. – **64**. – P. 91–97.
19. Запороцкая И. В., Чернозатонский Л. А. Исследование механизма положительного влияния фуллерена на процессы восстановления про-странственной памяти // *Вестник новых медицинских технологий*. – 2005. – Т.2. – С. 117–119.
20. Yang D., Wang M., Chen I. et al. Systemic administration of a water-soluble hexasulfonated C(60) (FC(4)S) reduces cerebral ischemia-induced infarct volume in gerbils // *Neurosci. Lett.* – 2001. – **311**, №2. – P. 121-124.
21. Kim J., Lee M. Fullerene inhibits beta-amyloid peptide aggregation // *Biochem. Biophys. Res. Commun.* – 2003. – **303**, №2. – P. 576-579.
22. Marsagishvili L., Bobylev A., Shpagina M. et al. Effect of fullerenes C<sub>60</sub> on X-protein amyloids // *Biofizika*. – 2009. – **54**, № 2. – P. 202–205.
23. Bobylev A., Kornev A., Bobyleva L. et al. Fullerenolates: metallated polyhydroxylated fullerenes with potent anti-amyloid activity // *Org. Biomol. Chem.* – 2011. – **9**. – P. 5714-5519.
24. Ширинкин С. В. Медицинские нанотехнологии. Использование природных фуллеренов при патологии памяти, внимания и реакций // *Мир психологии*. – 2007. – **3**, № 51. – С. 170–174.
25. Andrievsky G., Zhmuro A., Zabobonina L. First clinical case of treatment of patient (volunteer) with rectal adenocarcinoma by hydrated fullerenes: natural course of the disease or non-specific anticancer activity // *The Electrochemical Society Interface, Canada: Spring*. – 2000. – P. 700.
26. Chem C., Xing G., Wang J. et al. Multihydroxylated (Gd@C82(OH)22)n nanoparticles: antineoplastic activity of high efficiency and low toxicity // *Nanoletters*. – 2005. – **5**. – P. 2050–2057.
27. Ryan J., Bateman H., Stover A. et al. Fullerene nanomaterials inhibit the allergic response // *J. Immunology*. – 2007. – **179**, № 1. – P. 665–672.
28. Huang Y., Shen C., Luh T. et al. Blokade of apoptotic signaling of transforming growth factor-beta in human hepatoma cells by carboxyfullerene // *Eur. J. Biochem.* – 1998. – **254**, № 1. – P. 38–43.
29. Lin Y., Lei H., Luh T. et al. Light-independent inactivation of dengue-2 virus by carboxyfullerene C3 isomer // *Virology*. – 2000. – **276**, №1. – P. 258–262.
30. Piotrovsky L., Dumpis M., Poznykova L. et al. Study of the biological activity of the adducts of fullerenes with poly(N-vinylpyrrolidone) // *Mol. Mat.* – 2000. – **13**. – P. 41–50.
31. Миллер Г. Г., Куц А. А., Романова В. С. Средство для ингибирования ВИЧ и ЦМВ-инфекций и способ их ингибирования / RU 2196602. – 2003. – С. 1.
32. Tsao N., Luh T., Chou C. et al. In vitro action of carboxyfullerene // *J. Antimicrob. Chemother.* – 2002. – **49**, № 4. – P. 641–649.
33. Bosi S., Da Ros T., Spalluto G. et al. A new multi- charged C<sub>60</sub> derivative: synthesis and biological properties // *Eur. J. Org. Chem.* – 2002. – **17**. – P. 2928–2934.
34. Kornev A., Peregudov A., Martynenko V. et al. Synthesis and antiviral activity of highly water-soluble polycarboxylic derivatives of [70]fullerene // *Chem. Commun.* – 2011. – **47**, № 29. – P. 8298-8300.
35. Chawla P., Chawla V., Maheswari R. et al. Fullerenes: From carbon to nanomedicine // *Mini Reviews in Medicinal Chem.* – 2010. – **10**, № 8. – P. 662-677.
36. Kornev A., Khakina E., Troyanov S. et al. Facile preparation of amine and amino acid adducts of [60]fullerene using chlorofullerene C60Cl6 as a precursor // *Chem. Commun.* – 2012. – **48**, № 44. – P. 5461-5463.
37. Ciccolini F. Identification of two distinct types of multipotent neural precursors that appear sequentially during CNS development // *Mol. Cell Neurosci.* – 2001. – **17**, № 5. – P. 895-907.
38. Sun Y Shi J., Lu P. Neurotrophic factors and neural stem cells (abstract) // Sheng Li Ke Xue Jin. Zhan. – 2002. – **33**, № 4. – P. 313 - 316.
39. Israsena N., Hu M., Fu W. et al. The presence of FGF2 signaling determines whether beta-catenin exerts effects on proliferation or neuronal differentiation of neural stem cells // *Dev. Biol.* – 2004. – **268**, № 1. – P. 220-231.
40. Andressen C., Stocker E., Klinz F. Nestin-Specific Green Fluorescent Protein Expression in Embryonic Stem Cell-Derived Neural Precursor Cell Used for Transplantation // *Stem Cell*. – 2001. – **19**, №5. – P. 419-424.
41. Caldwell M.A., He X., Wilkie N. et al. Growth factors regulate the survival and fate of cells derived from human neurospheres // *Nature Biotechnol.* – 2001. – **19**, № 5. – P. 475 – 480.
42. Bez A., Corsini E., Curti D. et al. Neurosphere and neurosphere-forming cells: morphological and ultrastructural characterization // *Brain Res.* – 2003. – **993**, № 1-2. – P. 18-29.
43. Campos L. Neurospheres: insights into neural stem cell biology // *J. Neurosci. Res.* – 2004. – **78**, № 6. – P. 761-769.